

What happens



.. a **X-ray source...**

INPUTS
~~Source photons+~~
~~Mirrors response+~~
~~Detector response+~~
~~All kinds of~~
~~Background s~~

OUTPUTS
 Images
 Light Curves
 Spectra



Take into account telescope response... and remaining bgds



...**mirrors, concentrators**

limitors on board
tes..



Remove "some" backgrounds and malfunctioning

INPUTS
 Source photons+
 Mirrors response+
 Detector response+
 All kinds of
 Background s

ectors
Microcal., etc.)



things to do

..since the birth of X-ray Astronomy in 1962, improvements were carried out in terms of sensitivity, angular resolution, energy resolution and energy bandpass




The Golden age of X-ray Astronomy



Chandra



XMM-Newton



Suzaku

Sensitivity:

$$S/N = S / (S+B)^{0.5}$$

$$\longrightarrow \propto t^{0.5}$$

**$S^{0.5}$ = Poisson Noise
source counts**

$$B_{\text{sky}} = \text{Const} \times \text{Sky region}$$

$$B_{\text{dark current}} = \text{Const} \times \text{det. reg.}$$

$$B_{\text{rea-out (electronic)}}^2 = \text{Const} \times \text{det. Reg.}$$

How to increase the sensitivity....



Increasing the collecting/effective Area



$$S = F \times A_{\text{eff}}$$



S/N increases.....

(...but sometime also the bgd increases)

the ESA (XMM-Newton) way

Reducing the B.




S/N increases



the NASA (Chandra) way...

High Resolution Mirror Assembly (HRMA): On-axis PSF

Radius encompassing NN% of the counts
as a function of the energy





Encircled energy vs. radius
at different energies




On-axis PSF size and shape

High Resolution Mirror Assembly (HRMA): Off-axis PSF




CDF-N 2Ms exposure

Resulting image on the focal plane of ACIS




XMM-Newton: the EPIC on-axis PSF



Mirror module	2	3	4
Instr. chain ^a	pn	MOS-1+RGS-1	MOS-2+RGS-2
	orbit/ground	orbit/ground	orbit/ground
<i>FWHM</i> ["]	< 12.5 ^b /6.6	4.3/6.0	4.4/4.5
<i>HEW</i> ["]	15.2/15.1	13.8/13.6	13.0/12.8


PSF FWHM higher than in *Chandra* but much larger effective area
Background (and confusion limit) can be an issue

XMM-Newton: the EPIC on-axis PSF




Encircled energy vs. radius
at different energies for the MOS1-2

Encircled energy vs. radius
at different energies for the pn




XMM-Newton: the EPIC off-axis PSF




90% radius (radius encompassing 90% of the incoming photons) vs. off-axis angle for the MOS1-2 at different energies

90% radius vs. off-axis angle for the pn at different energies




Chandra = angular resolution



Only four, robust shells
High-quality of shell production
to allow $< \text{arcsec}$ on-axis angular
resolution (the best so far in X-rays)

$$\vartheta_{crit} \propto \frac{\sqrt{\rho}}{E}$$

High Resolution Mirror Assembly (HRMA)



Ottica Wolter Type-I

Mirror diameters:
1.23, 0.99, 0.87 0.65 m


Mirror lengths: 84 cm

HRMA mass: 1500 kg

Focal length: 10 m


PSF FWHM: 0.5"

Chandra focal-plane detectors: CCDs




XMM-Newton = large effective area

3 modules, 58 shells



$$\vartheta_{crit} \propto \frac{\sqrt{\rho}}{E}$$



XMM-Newton: all instruments at work simultaneously

xmm observatory system






Figure 3

Wolter I solution




30 arc min
diameter circles



EPIC MOS

7 CCDs each 10.9 x 10.9 arcminutes




EPIC pn


12 CCDs each 13.6 x 4.4 arcmin

Full ir


pn CCD, $\approx 50\%$ to the MOS1-2, the rest to the grating spectrometers (RGS)




Mirrors and Effective Area



Chandra: High Resolution Mirror Assembly (HRMA): Effective Area



Effective area vs. Energy




Effective area vs. off-axis angle at different energies


Effect of vignetting

$$\vartheta_{crit} \propto \frac{\sqrt{\rho}}{E}$$

Chandra: quantum efficiency




Chandra: effective area



Chandra: vignetting


Ratio of the off-axis vs. on-axis counts at different off-axis angles



Hard X-ray photons are more difficult to focus



→ Vignetting

XMM-Newton: mirror effective (geometric) area




$$\vartheta_{crit} \propto \frac{\sqrt{\rho}}{E}$$

XMM-Newton: quantum efficiency




Strong decrease in the QE above 10 keV, where also the effective area due to the mirrors has a significant decrease

XMM-Newton: effective area




XMM-Newton: vignetting




Strong vignetting (as expected) for high-energy photons, partly compensated by the large effective area (e.g., wrt. *Chandra*)

**You will account for all this information
creating a file named
arf (ancillary response file)**

Chandra





XMM-Newton



$$\vartheta_{crit} \propto \frac{\sqrt{\rho}}{E}$$

Mirrors and PSF



**Intrinsic limit ($\theta = 1.22 \lambda/D$)
+ operations...**

Optical band=seeing

**X-rays= mirrors properties
+ mirror array assembly**


Point Spread Function (PSF) – describes the response of an imaging system to a point source or point object.

HEW (PSF), FWHM (PSF) = angular resolution



PSF = function of (x,y) or (r, θ) .

High Resolution Mirror Assembly (HRMA): On-axis PSF

Radius encompassing NN% of the counts
as a function of the energy





Encircled energy vs. radius
at different energies




On-axis PSF size and shape

High Resolution Mirror Assembly (HRMA): Off-axis PSF




CDF-N 2Ms exposure

Resulting image on the focal plane of ACIS




XMM-Newton: the EPIC on-axis PSF



Mirror module	2	3	4
Instr. chain ^a	pn	MOS-1+RGS-1	MOS-2+RGS-2
	orbit/ground	orbit/ground	orbit/ground
<i>FWHM</i> ["]	< 12.5 ^b /6.6	4.3/6.0	4.4/4.5
<i>HEW</i> ["]	15.2/15.1	13.8/13.6	13.0/12.8


PSF FWHM higher than in *Chandra* but much larger effective area
Background (and confusion limit) can be an issue

XMM-Newton: the EPIC on-axis PSF




Encircled energy vs. radius at different energies for the MOS1-2

Encircled energy vs. radius at different energies for the pn




XMM-Newton: the EPIC off-axis PSF




90% radius (radius encompassing 90% of the incoming photons) vs. off-axis angle for the MOS1-2 at different energies

90% radius vs. off-axis angle for the pn at different energies



XMM-Newton: effective area dependence on the filter choice




To avoid contamination from bright, soft objects (e.g., stars), a medium/thick filter is adopted

Last but not least....


Energy resolution

Chandra: energy resolution



Typical CCD resolution
100-150 eV

XMM-Newton: energy resolution



$$\Delta E(\text{FWHM})/E \propto E^{-1/2} \quad (E \text{ in keV})$$

**You will account for all this information
creating a file named
rmf (redistribution matrix file)**